

# KDM5A Rabbit pAb

Catalog No.: A7238

1 Publications

## Basic Information

### Observed MW

192kDa

### Calculated MW

192kDa

### Category

Primary antibody

### Applications

ELISA, WB, IP, ChIP

### Cross-Reactivity

Human, Rat

## Background

This gene encodes a member of the Jumonji, AT-rich interactive domain 1 (JARID1) histone demethylase protein family. The encoded protein plays a role in gene regulation through the histone code by specifically demethylating lysine 4 of histone H3. The encoded protein interacts with many other proteins, including retinoblastoma protein, and is implicated in the transcriptional regulation of Hox genes and cytokines. This gene may play a role in tumor progression.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

**ChIP** 5µg antibody for  
10µg-15µg of Chromatin

## Immunogen Information

### Gene ID

5927

### Swiss Prot

P29375

### Immunogen

Recombinant protein of human KDM5A

### Synonyms

RBP2; RBBP2; RBBP-2; KDM5A

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Validation Data



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg KDM5A antibody (A7238). Western blot was performed from the immunoprecipitate using KDM5A antibody (A7238) at a dilution of 1:500.



Western blot analysis of various lysates using KDM5A Rabbit pAb (A7238) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 60s.



Chromatin immunoprecipitation analysis of extracts of HCT116 cells, using KDM5A antibody (A7238) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.