

# Acetyl-Histone H3-K27 Rabbit pAb

Catalog No.: A7253

64 Publications

## Basic Information

### Observed MW

17kDa

### Calculated MW

15kDa

### Category

Primary antibody

### Applications

ELISA, WB, IHC-P, IF/ICC, IP, ChIP, ChIP-seq

### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

## Recommended Dilutions

**WB** 1:1000 - 1:5000**IHC-P** 1:50 - 1:200**IF/ICC** 1:50 - 1:200**IP** 0.5ug-4ug antibody for  
200ug-400ug extracts of  
whole cells**ChIP** 5μg antibody for  
5μg-10μg of Chromatin**ChIP-seq** 1:20 - 1:100

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Histone H3 (NP\_003520.1).

### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; Acetyl-Histone H3-K27

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Validation Data



Western blot analysis of lysates from NIH/3T3 cells, using Acetyl-Histone H3-K27 Rabbit pAb (A7253) at 1:2000 dilution. NIH/3T3 cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.




Western blot analysis of lysates from C6 cells, using Acetyl-Histone H3-K27 Rabbit pAb (A7253) at 1:2000 dilution. C6 cells were treated by TSA (1  $\mu$ M) at 37°C for 18 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.




Chromatin immunoprecipitation was performed with cross-linked chromatin from 293T, using Acetyl-Histone H3-K27 Rabbit pAb antibody (A7253) and rabbit IgG (AC005). The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram constructed the ratios of the ratio of the immunoprecipitated DNA versus the input.


## Validation Data




Immunofluorescence analysis of C6 cells treated by TSA (upper left) and untreated C6 cells (upper right) using Acetyl-Histone H3-K27 Rabbit pAb (red, A7253) at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH-3T3 cells treated by TSA (upper left) and untreated NIH-3T3 cells (upper right) using Acetyl-Histone H3-K27 Rabbit pAb (red, A7253) at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of U-2 OS cells treated by TSA (upper left) and untreated U-2 OS cells (upper right) using Acetyl-Histone H3-K27 Rabbit pAb (red, A7253) at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded Human colon using Acetyl-Histone H3-K27 Rabbit pAb (A7253) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Human spleen using Acetyl-Histone H3-K27 Rabbit pAb (A7253) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Mouse lung using Acetyl-Histone H3-K27 Rabbit pAb (A7253) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Rat lung using Acetyl-Histone H3-K27 Rabbit pAb (A7253) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.