

# AKT1S1 Knockdown HeLa Cell Lysate, Heterozygous

Catalog No.: RM02045

### **Basic Information**

Catalog No.

RM02045

Category

Cell Lysate

**Parental Cell line** 

HeLa

Genotype

Knockdown

### **Background**

AKT1S1 is a proline-rich substrate of AKT (MIM 164730) that binds 14-3-3 protein (see YWHAH, MIM 113508) when phosphorylated (Kovacina et al., 2003 [PubMed 12524439]).[supplied by OMIM, Mar 2008]

#### **Gene Information**

### **Gene Symbol**

AKT1S1

### **Species**

Human

### Gene ID

84335

#### **Swiss Prot**

Q96B36

### Synonyms

Lobe; PRAS40

### **Contact**

8		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

### **Product Information**

### **Description**

AKT1S1 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:47bp deletion in exon2

Allele-2:48bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions** 

**Amount** 

4°C

50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20  $^{\circ}$ C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

WT CTATGCTGCCCATG\*\*\*\*\*\*\*\*GCACTGGCCCACAG
Mut CTATGCTGCCCATG\*\*\*Deletion\*\*\*GCACTGGCCCACAG
Allele-1: 47bp deletion in exon2

WT CTATGCTGCCCATG\*\*\*\*\*\*\*\*\*\*\*\*\*CACTGGCCCACAGG
Mut CTATGCTGCCCATG\*\*\*Deletion\*\*\*CACTGGCCCACAGG

Allele-2: 48bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and AKT1S1 Knockdown (KD) HeLa cells, using sanger sequencing.