

Recombinant Mouse IL-4 Protein

Catalog No.: RP01161 **Recombinant** **1 Publications**

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	16189	P07750

Tags

C-His

Synonyms

IL-4; BSF-1; IL4

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Interleukin-4, also known as IL4, is a secreted protein that belongs to the IL-4 / IL-13 family. Interleukin-4 / IL4 has many biological roles, including the stimulation of activated B-cell and T-cell proliferation. It enhances both secretion and cell surface expression of IgE and IgG1. Interleukin-4 / IL4 also regulates the expression of the low-affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. Interleukin-4 is essential for the switching of B cells to IgE antibody production and the maturation of T helper (Th) cells toward the Th2 phenotype.

Basic Information

Description

Recombinant Mouse IL-4 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (His23-Ser140) of mouse IL-4 (Accession #NP_067258.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

1. Measured in a cell proliferation assay using MC/9 2 mouse mast cells. The ED₅₀ for this effect is 2.8-11.3 pg/mL, corresponding to a specific activity of 8.85×10⁷~3.57×10⁸ units/mg. 2. Measured in a cell proliferation assay using HT-2 mouse T cells. The ED₅₀ for this effect is 0.21-0.85 ng/mL, corresponding to a specific activity of 1.18×10⁶~4.76×10⁶ units/mg.

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.


After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn


Validation Data




Recombinant Mouse IL-4 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 20 kDa.




Recombinant Mouse IL-4 promote the proliferation of MC/9 2 mouse mast cells. The ED₅₀ for this effect is 2.8-11.3 pg/mL, corresponding to a specific activity of $8.85 \times 10^7 \sim 3.57 \times 10^8$ units/mg.




Recombinant Mouse IL4 stimulates cell proliferation of the HT-2 mouse T cells. The ED₅₀ for this effect is 0.21-0.85 ng/mL, corresponding to a specific activity of $1.18 \times 10^6 \sim 4.76 \times 10^6$ units/mg.



Active Recombinant Mouse IL-4 induces Mouse Bone marrow derived macrophage (BMDM) differentiation into M2-type. 20 ng/mL of Recombinant Mouse IL-4 and Mouse IL-13 can effectively upregulate the expression of M2-type biomarker Arg-1. (Customer feedback data)



Mouse IL-4 (10 ng/mL) and GM-CSF (20 ng/mL) induce C57BL/6J Mouse cells differentiate into BMDC. After 6 days, up to 75% of cells in the system expressed DC biomarker CD11c on cell surface, among which more than 15% cells have high expression of MHCII, shows that the induction was successful. (Customer feedback data)



Recombinant Mouse IL-4 (10 ng/mL) and GM-CSF protein (20 ng/mL) induce C57BL/6J Mouse bone marrow cells to differentiate into BMDC. After 7 days, up to 89.4% of cells in the system expressed DC biomarker CD11c on cell surface, among which more than half of the cells have high expression of CD80 and CD86 proteins. (Customer feedback data)