

MMP14/MT1-MMP Rabbit mAb

Catalog No.: A0067 **Recombinant** **3 Publications**

Basic Information

Observed MW

52kDa/60kDa

Calculated MW

66kDa

Category

Primary antibody

Applications

ELISA, WB, FC, IHC-P

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0211

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the protein encoded by this gene is a member of the membrane-type MMP (MT-MMP) subfamily; each member of this subfamily contains a potential transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. This protein activates MMP2 protein, and this activity may be involved in tumor invasion.

Recommended Dilutions

WB	1:500 - 1:2000
FC	1:100 - 1:500
IHC-P	1:50 - 1:200

Immunogen Information

Gene ID

4323

Swiss Prot

P50281

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human MMP14/MMP14/MT1-MMP (P50281).

Synonyms

MMP-14; MMP-X1; MT-MMP; MT1MMP; MTMMP1; WNCHRS; MT1-MMP; MT-MMP 1; MMP14/MT1-MMP

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification




Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data



Flow cytometry: 1X10⁶ MCF7 cells (negative control, left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067, 2 µg/mL, orange line) or ABflo® 488 Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by Alexa Fluor® 488 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).




Flow cytometry: 1X10⁶ MCF7 cells (negative control, left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067, 2 µg/mL, orange line) or ABflo® 488 Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by Alexa Fluor® 488 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10⁶ MCF7 cells (negative control, left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067, 2 µg/mL, orange line) or ABflo® 647 Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).

Validation Data



Flow cytometry: 1×10^6 MCF7 cells (negative control, left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067, 2 $\mu\text{g}/\text{mL}$, orange line) or ABflo® 647 Rabbit IgG isotype control (AC042, 2 $\mu\text{g}/\text{mL}$, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).