Cytokeratin 13 (KRT13) Rabbit mAb

www.abclonal.com

ABclonal

Catalog No.: A0411 Recombinant 5 Publications

Basic Information

Observed MW

50 kDa

Calculated MW

50 kDa

Category

Primary antibody

Applications

WB,IF-P,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1824

Background

The protein encoded by this gene is a member of the keratin gene family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Most of the type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. This type I cytokeratin is paired with keratin 4 and expressed in the suprabasal layers of non-cornified stratified epithelia. Mutations in this gene and keratin 4 have been associated with the autosomal dominant disorder White Sponge Nevus. The type I cytokeratins are clustered in a region of chromosome 17q21.2. Alternative splicing of this gene results in multiple transcript variants; however, not all variants have been described.

Recommended Dilutions

WB 1:5000 - 1:20000

1:100 - 1:1000 IF-P

IHC-P 1:500 - 1:2000

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on vour specific assav requirements.

Immunogen Information

Gene ID Swiss Prot 3860 P13646

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

K13; CK13; WSN2; Cytokeratin 13 (KRT13)

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	1	www.abclonal.com.cn


Product Information

Source Isotype **Purification** Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using Cytokeratin 13 (KRT13) (KRT13) Rabbit mAb (A0411) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 1s.




Immunofluorescence analysis of paraffinembedded rat bladder using Cytokeratin 13 (KRT13) (KRT13) Rabbit mAb (A0411) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffinembedded Human esophagus tissue using Cytokeratin 13 (KRT13) Rabbit mAb (A0411) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunofluorescence analysis of paraffinembedded human cervix cancer using Cytokeratin 13 (KRT13) (KRT13) Rabbit mAb (A0411) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffinembedded Human tonsil tissue using Cytokeratin 13 (KRT13) Rabbit mAb (A0411) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunofluorescence analysis of paraffinembedded mouse bladder using Cytokeratin 13 (KRT13) (KRT13) Rabbit mAb (A0411) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.