

[KO Validated] Hexokinase II Rabbit pAb

Catalog No.: A0994SP

KO Validated

53 Publications

Basic Information

Observed MW

102 kDa

Calculated MW

102 kDa

Category

Primary antibody

Applications

WB, IP, IF/ICC, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate of glycolysis seen in rapidly growing cancer cells.

Recommended Dilutions

WB 1:5000- 1:15000**IP** 0.5 µg - 4 µg antibody for
200 µg - 400 µg extracts of
whole cells**IF/ICC** 1:100 - 1:400**IHC-P** 1:200 - 1:800**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on your
specific assay
requirements. For high-ratio
antibody dilutions
(≥1:10000) a sequential
dilution method is strongly
recommended to ensure
measurement accuracy.

Immunogen Information

Gene ID

3099

Swiss Prot

P52789

Immunogen

This information is considered to be commercially sensitive.

Synonyms

HKII; HXK2; Hexokinase II

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide.

May contain 0.05% BSA as specified on the Certificate of Analysis.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Western blot analysis of lysates from Mouse heart using [KO Validated] Hexokinase II Rabbit pAb (A0994SP) at 1:10000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3 s.




Western blot analysis of various lysates using [KO Validated] Hexokinase II Rabbit pAb (A0994SP) at 1:10000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10 s.




Western blot analysis of lysates from wild type (WT) and Hexokinase II knockout (KO) HeLa cells using [KO Validated] Hexokinase II Rabbit pAb (A0994SP) at 1:10000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10 s.

Validation Data



Confocal imaging of HeLa cells using [KO Validated] Hexokinase II Rabbit pAb (A0994SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunohistochemistry analysis of paraffin-embedded Human colon tissue using [KO Validated] Hexokinase II Rabbit pAb (A0994SP) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.