

ZSCAN4C Rabbit pAb

Catalog No.: A10205

1 Publications

Basic Information

Observed MW

57kDa

Calculated MW

57kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Mouse

Background

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Acts upstream of or within negative regulation of mitotic recombination; regulation of transcription by RNA polymerase II; and telomere maintenance via telomere lengthening. Located in chromosome, telomeric region; cytoplasm; and nucleus. Is expressed in 1-cell stage embryo; 2-cell stage embryo; primary oocyte; and secondary oocyte. Orthologous to human ZSCAN4 (zinc finger and SCAN domain containing 4).

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

245109

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Gm397; Zscan4d; XM_142517; ZSCAN4C

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification

Storage

Store at -20°C

Validation Data



Western blot analysis of lysates from mouse kidney, using ZSCAN4C Rabbit pAb (A10205) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.