BCL2A1 Rabbit pAb

Catalog No.: A1105 1 Publications



Basic Information

Observed MW

Refer to figures

Calculated MW

20kDa

Category

Primary antibody

Applications

WB

Cross-Reactivity

Human

Background

This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and pro-apoptotic regulators that are involved in a wide variety of cellular activities such as embryonic development, homeostasis and tumorigenesis. The protein encoded by this gene is able to reduce the release of pro-apoptotic cytochrome c from mitochondria and block caspase activation. This gene is a direct transcription target of NF-kappa B in response to inflammatory mediators, and is upregulated by different extracellular signals, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), CD40, phorbol ester and inflammatory cytokine TNF and IL-1, which suggests a cytoprotective function that is essential for lymphocyte activation as well as cell survival. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID

Swiss Prot

Q16548

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GRS; ACC1; ACC2; BFL1; ACC-1; ACC-2; HBPA1; BCL2L5; BCL2A1

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.