

TRADD Rabbit pAb

Catalog No.: A1145

3 Publications

Basic Information

Observed MW

36kDa

Calculated MW

34kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse

Background

Predicted to enable death domain binding activity; identical protein binding activity; and signaling receptor binding activity. Involved in positive regulation of hair follicle development. Acts upstream of or within extrinsic apoptotic signaling pathway. Located in cytoplasm. Is expressed in adrenal gland; genitourinary system; liver; lung; and spleen. Orthologous to human TRADD (TNFRSF1A associated via death domain).

Recommended Dilutions

WB 1:1000 - 1:5000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

8717

Swiss Prot

Q15628

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

9130005N23Rik; TRADD

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from MCF7 cells, using TRADD Rabbit pAb (A1145) at 1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.