# SOX30 Rabbit pAb

Catalog No.: A11715 2 Publications



### **Basic Information**

### **Observed MW**

90kDa

#### **Calculated MW**

82kDa

### Category

Primary antibody

### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. The encoded protein acts as a transcriptional regulator when present in a complex with other proteins. It can activate p53 transcription to promote tumor cell apoptosis in lung cancer. The protein may be involved in the differentiation of developing male germ cells. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been identified on chromosome 5.

### **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$ 

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 11063 094993

#### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

SOX30

### **Contact**

| <b>a</b>          |   | 400-999-6126              |
|-------------------|---|---------------------------|
| $\bowtie$         |   | cn.market@abclonal.com.cn |
| $\overline{\Box}$ | Т | www.ahclonal.com.cn       |


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using SOX30 Rabbit pAb (A11715) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.