RUNX2 Rabbit mAb

Catalog No.: A11753 Recombinant 14 Publications



Basic Information

Observed MW

60kDa

Calculated MW

57kDa

Category

Primary antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC0362

Background

This gene is a member of the RUNX family of transcription factors and encodes a nuclear protein with an Runt DNA-binding domain. This protein is essential for osteoblastic differentiation and skeletal morphogenesis and acts as a scaffold for nucleic acids and regulatory factors involved in skeletal gene expression. The protein can bind DNA both as a monomer or, with more affinity, as a subunit of a heterodimeric complex. Two regions of potential trinucleotide repeat expansions are present in the N-terminal region of the encoded protein, and these and other mutations in this gene have been associated with the bone development disorder cleidocranial dysplasia (CCD). Transcript variants that encode different protein isoforms result from the use of alternate promoters as well as alternate splicing.

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5μg-4μg antibody for 400μg-600μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID Swiss Prot860
Q13950

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CCD; AML3; CCD1; CLCD; OSF2; CBFA1; OSF-2; PEA2aA; PEBP2aA; CBF-alpha-1; RUNX2

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Immunoprecipitation analysis of 300 μ g extracts of Mouse testis cells using 3 μ g RUNX2 antibody (A11753). Western blot was performed from the immunoprecipitate using RUNX2 antibody (A11753) at a dilution of 1:500.




Western blot analysis of various lysates using RUNX2 Rabbit mAb (A11753) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Western blot analysis of various lysates using RUNX2 Rabbit mAb (A11753) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.