# STIP1 Rabbit pAb

Catalog No.: A1219 2 Publications



## **Basic Information**

#### **Observed MW**

63kDa

#### **Calculated MW**

63kDa

#### Category

Primary antibody

#### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

STIP1 is an adaptor protein that coordinates the functions of HSP70 (see HSPA1A; MIM 140550) and HSP90 (see HSP90AA1; MIM 140571) in protein folding. It is thought to assist in the transfer of proteins from HSP70 to HSP90 by binding both HSP90 and substrate-bound HSP70. STIP1 also stimulates the ATPase activity of HSP70 and inhibits the ATPase activity of HSP90, suggesting that it regulates both the conformations and ATPase cycles of these chaperones (Song and Masison, 2005 [PubMed 16100115]).

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

IHC-P 1:50 - 1:200

**ELISA** Recommended starting concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID**Swiss Prot
10963
P31948

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

HOP; P60; STI1; STI1L; HEL-S-94n; IEF-SSP-3521; STIP1

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification


# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using STIP1 Rabbit pAb (A1219) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunohistochemistry analysis of paraffinembedded Rat testis using STIP1 Rabbit pAb (A1219) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.