IL10 Rabbit mAb

Catalog No.: A12255 Recombinant 24 Publications



Basic Information

Observed MW

17kDa

Calculated MW

21kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Rat

CloneNo number

ARC0650

Background

The protein encoded by this gene is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Recommended Dilutions

WB 1:1000 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID3586

Swiss Prot
P22301

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

CSIF; TGIF; GVHDS; IL-10; IL10A; IL10

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using IL10 Rabbit mAb (A12255) at 1:1000 dilution incubated overnight at 4° C. THP-1 cells were treated with LPS (100ng/ml) and Brefeldin A (300ng/ml) for different times of stimulation.


Secondary antibody: HRP-conjugated Goat anti-Rabbit $IgG\ (H+L)\ (AS014)$ at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.




Western blot analysis of lysates from Rat thymus, using IL10 Rabbit mAb (A12255) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Confocal imaging of HeLa cells using IL10 Rabbit mAb (A12255,dilution 1:100)(Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012,dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.