# **SCD5 Rabbit pAb**

Catalog No.: A13127 2 Publications



## **Basic Information**

#### **Observed MW**

38kDa

#### **Calculated MW**

38kDa

#### Category

Primary antibody

### **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human

## **Background**

Stearoyl-CoA desaturase (SCD; EC 1.14.99.5) is an integral membrane protein of the endoplasmic reticulum that catalyzes the formation of monounsaturated fatty acids from saturated fatty acids. SCD may be a key regulator of energy metabolism with a role in obesity and dislipidemia. Four SCD isoforms, Scd1 through Scd4, have been identified in mouse. In contrast, only 2 SCD isoforms, SCD1 (MIM 604031) and SCD5, have been identified in human. SCD1 shares about 85% amino acid identity with all 4 mouse SCD isoforms, as well as with rat Scd1 and Scd2. In contrast, SCD5 shares limited homology with the rodent SCDs and appears to be unique to primates (Wang et al., 2005 [PubMed 15907797]).

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:100 - 1:500

**ELISA** Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

 Gene ID
 Swiss Prot

 79966
 Q865K9

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

SCD2; SCD4; ACOD4; FADS4; HSCD5; DFNA79; SCD5

## **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using SCD5 Rabbit pAb (A13127) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.



Immunofluorescence analysis of Neuro-2a cells using SCD5 Rabbit pAb (A13127) at a dilution of 1:300 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.