

FAH Rabbit pAb

Catalog No.: A13492 **4 Publications**

Basic Information

Observed MW

40-45kDa

Calculated MW

46kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Predicted to enable fumarylacetoacetase activity. Predicted to be involved in L-phenylalanine catabolic process; homogentisate catabolic process; and tyrosine catabolic process. Predicted to act upstream of or within arginine catabolic process. Located in extracellular exosome. Implicated in tyrosinemia type I.

Recommended Dilutions

WB 1:2000 - 1:8000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

2184

Swiss Prot

P16930

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

FAH

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates, using FAH Rabbit pAb (A13492) at 1:7000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.