## **HDAC4** Rabbit mAb

Catalog No.: A13510 Recombinant 3 Publications



## **Basic Information**

#### **Observed MW**

140kDa

### **Calculated MW**

119kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human

#### CloneNo number

ARC0714

## **Background**

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**9759

Swiss Prot
P56524

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

HD4; AHO3; BDMR; HDACA; HA6116; HDAC-4; HDAC-A; NEDCHF; NEDCHID; HDAC4

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{a}$	ī	www.ahclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates, using HDAC4 Rabbit mAb (A13510) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.