# Collagen I/COL1A1 Rabbit pAb

Catalog No.: A1352 149 Publications



## **Basic Information**

## **Observed MW**

220 kDa

#### **Calculated MW**

139 kDa

## Category

Primary antibody

## **Applications**

WB,IHC-P,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes the pro-alpha1 chains of type I collagen whose triple helix comprises two alpha1 chains and one alpha2 chain. Type I is a fibril-forming collagen found in most connective tissues and is abundant in bone, cornea, dermis and tendon. Mutations in this gene are associated with osteogenesis imperfecta types I-IV, Ehlers-Danlos syndrome type VIIA, Ehlers-Danlos syndrome Classical type, Caffey Disease and idiopathic osteoporosis. Reciprocal translocations between chromosomes 17 and 22, where this gene and the gene for platelet-derived growth factor beta are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans, resulting from unregulated expression of the growth factor. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene.

## **Recommended Dilutions**

**WB** 1:1000 - 1:6000

IHC-P 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

## Immunogen Information

 Gene ID
 Swiss Prot

 1277
 P02452

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

OI1; OI2; OI3; OI4; EDSC; CAFYD; EDSARTH1; Collagen I/COL1A1

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	1	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using Collagen I/COL1A1 Rabbit pAb (A1352) at 1:5000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Negative control (NC): HT-29.

Exposure time: 30s.



Immunohistochemistry analysis of paraffinembedded Human cervix tissue using Collagen I/COL1A1 Rabbit pAb (A1352) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of NIH/3T3 cells using Collagen I/COL1A1 Rabbit pAb (A1352) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.