

TNFAIP8L2 Rabbit pAb

Catalog No.: A13698 **1 Publications**

Basic Information

Observed MW

18kDa

Calculated MW

21kDa

Category

Primary antibody

Applications

WB,IF/ICC,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Predicted to be involved in negative regulation of T cell activation and negative regulation of inflammatory response. Predicted to be active in cytoplasm.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
25µg-100µg extracts of
whole cells

ELISA Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

79626

Swiss Prot

Q6P589

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TIPE2; TNFAIP8L2

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using TNFAIP8L2 Rabbit pAb (A13698) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.




Immunoprecipitation analysis of 25 µg extracts of Mouse spleen cells using 3 µg TNFAIP8L2 antibody (A13698). Western blot was performed from the immunoprecipitate using TNFAIP8L2 antibody (A13698) at a dilution of 1:1000.



Immunofluorescence analysis of C6 cells using TNFAIP8L2 Rabbit pAb (A13698) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using TNFAIP8L2 Rabbit pAb (A13698) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using TNFAIP8L2 Rabbit pAb (A13698) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.