

IFIT2 Rabbit pAb

Catalog No.: A13760 **1 Publications**

Basic Information

Observed MW

50kDa

Calculated MW

55kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human

Background

Enables RNA binding activity. Involved in negative regulation of protein binding activity; positive regulation of apoptotic process; and response to virus. Located in endoplasmic reticulum.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

3433

Swiss Prot

P09913

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

P54; G10P2; IFI54; ISG54; cig42; IFI-54; IFIT-2; ISG-54; GARG-39; IFI-54K; ISG-54K; ISG-54 K; IFIT2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using IFIT2 Rabbit pAb (A13760) at 1:3000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.