

HP1BP3 Rabbit pAb

Catalog No.: A14209

Basic Information

Observed MW

75kDa

Calculated MW

61kDa

Category

Primary antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables DNA binding activity and nucleosome binding activity. Involved in several processes, including cellular response to hypoxia; heterochromatin organization; and regulation of nucleus size. Located in chromosome and nuclear speck.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

50809

Swiss Prot

Q5SSJ5

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

HP1BP74; HP1-BP74; HP1BP3

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification


Storage

Store at -20°C


Validation Data




Western blot analysis of lysates from LO2 cells, using HP1BP3 Rabbit pAb (A14209) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using HP1BP3 Rabbit pAb (A14209) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using HP1BP3 Rabbit pAb (A14209) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using HP1BP3 Rabbit pAb (A14209) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.