FTO Rabbit pAb

Catalog No.: A1438 25 Publications



Basic Information

Observed MW

60kDa

Calculated MW

58kDa

Category

Primary antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene is a nuclear protein of the AlkB related non-haem iron and 2-oxoglutarate-dependent oxygenase superfamily but the exact physiological function of this gene is not known. Other non-heme iron enzymes function to reverse alkylated DNA and RNA damage by oxidative demethylation. Studies in mice and humans indicate a role in nervous and cardiovascular systems and a strong association with body mass index, obesity risk, and type 2 diabetes.

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene IDSwiss Prot
79068
Q9C0B1

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GDFD; ALKBH9; BMIQ14; FTO

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using FTO Rabbit pAb (A1438) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunoprecipitation analysis of 300 μ g extracts of SH-SY5Y cells using 3 μ g FTO antibody (A1438). Western blot was performed from the immunoprecipitate using FTO antibody (A1438) at a dilution of