# SAA1 Rabbit pAb

Catalog No.: A14553 2 Publications



# **Basic Information**

#### **Observed MW**

16kDa

### **Calculated MW**

12kDa

#### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Mouse, Rat

# **Background**

This gene encodes a member of the serum amyloid A family of apolipoproteins. The encoded preproprotein is proteolytically processed to generate the mature protein. This protein is a major acute phase protein that is highly expressed in response to inflammation and tissue injury. This protein also plays an important role in HDL metabolism and cholesterol homeostasis. High levels of this protein are associated with chronic inflammatory diseases including atherosclerosis, rheumatoid arthritis, Alzheimer's disease and Crohn's disease. This protein may also be a potential biomarker for certain tumors. Finally, antimicrobial activity against S. aureus and E. coli resides in the N-terminal portion of the mature protein. Alternate splicing results in multiple transcript variants that encode the same protein. A pseudogene of this gene is found on chromosome 11.

### **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**6288
Swiss Prot
PODJI8

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

SAA; PIG4; SAA2; TP53I4; SAA1

### **Contact**

| <u>a</u>  | 400-999-6126              |
|-----------|---------------------------|
| $\bowtie$ | cn.market@abclonal.com.cn |
|           | www.abclonal.com.cn       |


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using SAA1 Rabbit pAb (A14553) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.