

CDK6 Rabbit pAb

Catalog No.: A1545SP

14 Publications

Basic Information

Observed MW

37 kDa

Calculated MW

37 kDa

Category

Primary antibody

Applications

WB, IP, ELISA

Cross-Reactivity

Human

Background

The protein encoded by this gene is a member of the CMGC family of serine/threonine protein kinases. This kinase is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression and G1/S transition. The activity of this kinase first appears in mid-G1 phase, which is controlled by the regulatory subunits including D-type cyclins and members of INK4 family of CDK inhibitors. This kinase, as well as CDK4, has been shown to phosphorylate, and thus regulate the activity of, tumor suppressor protein Rb. Altered expression of this gene has been observed in multiple human cancers. A mutation in this gene resulting in reduced cell proliferation, and impaired cell motility and polarity, and has been identified in patients with primary microcephaly.

Recommended Dilutions

WB 1:3000 - 1:10000**IP** 0.5 µg - 4 µg antibody for
200 µg - 400 µg extracts of
whole cells**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on your
specific assay
requirements. For high-ratio
antibody dilutions
(≥1:10000)

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Immunogen Information

Gene ID

1021

Swiss Prot

Q00534

Immunogen

This information is considered to be commercially sensitive.

Synonyms

MCPH12; PLSTIRE; CDK6

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide.

May contain 0.05% BSA as specified on the Certificate of Analysis.

Validation Data



Immunoprecipitation of CDK6 from 200 µg extracts of Jurkat cells was performed using 1 µg of CDK6 Rabbit pAb (A1545SP). Rabbit IgG isotype control (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using CDK6 Rabbit pAb (A1545SP) at a dilution of 1:5000.



Western blot analysis of lysates from Jurkat cells using CDK6 Rabbit pAb (A1545SP) at 1:5000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1 s.