Survivin Rabbit pAb

Catalog No.: A1551 24 Publications



Basic Information

Observed MW

16 kDa

Calculated MW

16 kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene is a member of the inhibitor of apoptosis (IAP) gene family, which encode negative regulatory proteins that prevent apoptotic cell death. IAP family members usually contain multiple baculovirus IAP repeat (BIR) domains, but this gene encodes proteins with only a single BIR domain. The encoded proteins also lack a C-terminus RING finger domain. Gene expression is high during fetal development and in most tumors, yet low in adult tissues. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

1:500 - 1:1000

concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

Recommended starting

For highratio
antibody
dilutions
(≥1:10000)[]

a sequential dilution method is strongly recommend ed to ensure measureme

nt accuracy.

Immunogen Information

Gene ID332
Swiss Prot
015392

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

API4; EPR-1; Survivin

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Contact

2	400-999-6126
\bowtie	cn.market@abclonal.com.cr
•	www.abclonal.com.cr



Western blot analysis of various lysates, using Survivin Rabbit pAb (A1551) at 1:400 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.