

TEAD2 Rabbit pAb

Catalog No.: A15594 **4 Publications**

Basic Information

Observed MW

51kDa

Calculated MW

49kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human

Background

Enables several functions, including DNA-binding transcription factor activity; disordered domain specific binding activity; and transcription coactivator binding activity. Involved in hippo signaling; positive regulation of transcription, DNA-templated; and protein-containing complex assembly. Located in cytosol and nucleoplasm. Part of TEAD-YAP complex.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

8463

Swiss Prot

Q15562

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ETF; TEF4; TEF-4; TEAD-2; TEAD2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from A-431 cells, using TEAD2 Rabbit pAb (A15594) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.