Leader in Biomolecular Solutions for Life Science

www.abclonal.com

p90Rsk/RSK1/RPS6KA1 Rabbit pAb

Catalog No.: A15718 6 Publications

Basic Information

Observed MW
90kDa

Calculated MW
83kDa

Category
Primary antibody

Applications
WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity
Human, Mouse, Rat

Background

This gene encodes a member of the RSK (ribosomal S6 kinase) family of serine/threonine kinases. This kinase contains 2 nonidentical kinase catalytic domains and phosphorylates various substrates, including members of the mitogen-activated kinase (MAPK) signalling pathway. The activity of this protein has been implicated in controlling cell growth and differentiation. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:100 - 1:200
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
6195

Swiss Prot
Q15418

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
RSK; HU-1; RSK1; p90Rsk; MAPKAPK1; MAPKAPK1A; p90Rsk/RSK1/RPS6KA1

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using p90Rsk/RSK1/RPS6KA1 Rabbit pAb (A15718) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Immunofluorescence analysis of L929 cells using p90Rsk/RSK1/RPS6KA1 Rabbit pAb (A15718) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using p90Rsk/RSK1/RPS6KA1 Rabbit pAb (A15718) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen using p90Rsk/RSK1/RPS6KA1 Rabbit pAb (A15718) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.