

IGHMBP2 Rabbit pAb

Catalog No.: A16053

Basic Information

Observed MW

109kDa

Calculated MW

109kDa

Category

Primary antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a helicase superfamily member that binds a specific DNA sequence from the immunoglobulin mu chain switch region. Mutations in this gene lead to spinal muscle atrophy with respiratory distress type 1.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:100

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

3508

Swiss Prot

P38935

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

HCSA; HMN6; CATF1; CMT2S; SMARD1; SMUBP2; ZFAND7; IGHMBP2

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG



Purification

Affinity purification


Storage

Store at -20°C


Validation Data




Immunohistochemistry analysis of paraffin-embedded Rat brain using IGHMBP2 Rabbit pAb (A16053) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using IGHMBP2 Rabbit pAb (A16053) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human mammary cancer using IGHMBP2 Rabbit pAb (A16053) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver using IGHMBP2 Rabbit pAb (A16053) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.