# SIAH2 Rabbit pAb

Catalog No.: A16211 1 Publications



### **Basic Information**

#### **Observed MW**

35kDa

### **Calculated MW**

35kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human

# **Background**

This gene encodes a protein that is a member of the seven in absentia homolog (SIAH) family. The protein is an E3 ligase and is involved in ubiquitination and proteasome-mediated degradation of specific proteins. The activity of this ubiquitin ligase has been implicated in regulating cellular response to hypoxia.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**6478

Swiss Prot

043255

#### **Immunogen**

 $Synthetic\ peptide.\ This\ information\ is\ considered\ to\ be\ commercially\ sensitive.$ 

### **Synonyms**

hSiah2; SIAH2

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{\triangle}$	ī	www.ahclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of lysates from SKOV3 cells, using SIAH2 Rabbit pAb (A16211) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 5min.