# MYF5 Rabbit pAb

Catalog No.: A16227 6 Publications



### **Basic Information**

#### **Observed MW**

28kDa

### **Calculated MW**

28kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

### **Background**

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Predicted to contribute to E-box binding activity. Predicted to be involved in several processes, including muscle cell fate commitment; positive regulation of cell differentiation; and skeletal muscle cell differentiation. Predicted to act upstream of or within several processes, including animal organ development; regulation of cell-matrix adhesion; and somitogenesis. Located in nucleoplasm.

### **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID Swiss Prot** 4617 P13349

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

EORVA; bHLHc2; MYF5

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn

### **Product Information**


**Purification** Source Isotype Rabbit IgG Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from Mouse testis, using MYF5 Rabbit pAb (A16227) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.