

ASC/TMS1 Rabbit pAb

Catalog No.: A16672 34 Publications

Basic Information

Observed MW

19-25kDa

Calculated MW

22kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes an adaptor protein that is composed of two protein-protein interaction domains: a N-terminal PYRIN-PAAD-DAPIN domain (PYD) and a C-terminal caspase-recruitment domain (CARD). The PYD and CARD domains are members of the six-helix bundle death domain-fold superfamily that mediates assembly of large signaling complexes in the inflammatory and apoptotic signaling pathways via the activation of caspase. In normal cells, this protein is localized to the cytoplasm; however, in cells undergoing apoptosis, it forms ball-like aggregates near the nuclear periphery. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:1000 - 1:6000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
IP	0.5μg-4μg antibody for 400μg-600μg extracts of whole cells
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

29108

Swiss Prot

Q9ULZ3

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ASC; TMS; TMS1; CARD5; TMS-1; ASC/TMS1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Validation Data




Immunoprecipitation analysis of 200 µg extracts of Rat lung using 3 µg ASC/TMS1 antibody (A16672). Western blot was performed from the immunoprecipitate using ASC/TMS1 antibody (A16672) at a dilution of 1:1000.




Western blot analysis of various lysates using ASC/TMS1 Rabbit pAb (A16672) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Human esophageal cancer using ASC/TMS1 Rabbit pAb (A16672) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human tonsil using ASC/TMS1 Rabbit pAb (A16672) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of C6 cells using ASC/TMS1 Rabbit pAb (A16672) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Validation Data



Immunofluorescence analysis of NIH-3T3 cells using ASC/TMS1 Rabbit pAb (A16672) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using ASC/TMS1 Rabbit pAb (A16672) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.