VGLL4 Rabbit pAb

Catalog No.: A17058 2 Publications



Basic Information

Observed MW

31kDa

Calculated MW

31kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Mouse, Rat

Background

Predicted to enable transcription coactivator binding activity. Involved in negative regulation of Wnt signaling pathway; negative regulation of cell growth; and negative regulation of hippo signaling. Predicted to be located in nucleus.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID9686

Swiss Prot
Q14135

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

VGL-4; VGLL4

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using VGLL4 Rabbit pAb (A17058) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.