MIXL1 Rabbit pAb

Catalog No.: A17223 4 Publications



Basic Information

Observed MW

23kDa

Calculated MW

25kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables RNA polymerase II-specific DNA-binding transcription factor binding activity and sequence-specific double-stranded DNA binding activity. Predicted to be involved in several processes, including endodermal cell differentiation; hematopoietic progenitor cell differentiation; and positive regulation of mesoderm development. Predicted to act upstream of or within cell migration involved in gastrulation and hemopoiesis. Located in nucleoplasm.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID83881

Swiss Prot
Q9H2W2

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

MIX; MIXL; MILD1; MIXL1

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\overline{a}	ī	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using MIXL1 Rabbit pAb (A17223) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5min.