# ETV5 Rabbit pAb

Catalog No.: A17927



### **Basic Information**

#### **Observed MW**

70kDa

#### **Calculated MW**

58kDa

#### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Enables DNA-binding transcription activator activity, RNA polymerase II-specific and RNA polymerase II transcription regulatory region sequence-specific DNA binding activity. Involved in cellular response to oxidative stress; negative regulation of transcription by RNA polymerase II; and positive regulation of transcription by RNA polymerase II. Located in nucleoplasm.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**2119

Swiss Prot
P41161

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

ERM; ETV5

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Ī	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from A-431 cells, using ETV5 Rabbit pAb (A17927) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.