# **GPR35 Rabbit pAb**

Catalog No.: A17953 1 Publications



## **Basic Information**

#### **Observed MW**

34kDa

### **Calculated MW**

34kDa

### Category

Primary antibody

## **Applications**

WB,ELISA

### **Cross-Reactivity**

Mouse

## **Background**

Enables C-X-C chemokine receptor activity. Involved in several processes, including chemokine-mediated signaling pathway; negative regulation of voltage-gated calcium channel activity; and positive regulation of cytosolic calcium ion concentration. Is integral component of plasma membrane.

## **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID Swiss Prot** 2859 Q9HC97

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

GPR35

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn


## **Product Information**

**Purification Source** Isotype Affinity purification Rabbit IgG

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using GPR35 Rabbit pAb (A17953) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.