

[KO Validated] CDKN2C/p18-INK4C Rabbit pAb

Catalog No.: A18054 **KO** Validated

Basic Information

Observed MW

18kDa

Calculated MW

18kDa

Category

Primary antibody

Applications

ELISA, WB, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase inhibitors. This protein has been shown to interact with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. Ectopic expression of this gene was shown to suppress the growth of human cells in a manner that appears to correlate with the presence of a wild-type RB1 function. Studies in the knockout mice suggested the roles of this gene in regulating spermatogenesis, as well as in suppressing tumorigenesis. Two alternatively spliced transcript variants of this gene, which encode an identical protein, have been reported.

Recommended Dilutions

WB 1:500 - 1:2000**IF/ICC** 1:50 - 1:100

Immunogen Information

Gene ID

1031

Swiss Prot

P42773

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-168 of human CDKN2C/p18-INK4C (NP_523240.1).

Synonyms

p18; INK4C; p18-INK4C; 4C

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and CDKN2C/p18-INK4C knockout (KO) HeLa cells, using [KO Validated] CDKN2C/p18-INK4C Rabbit pAb (A18054) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.