

ZNF444 Rabbit pAb

Catalog No.: A18470

Basic Information

Observed MW

38kDa

Calculated MW

35kDa

Category

Primary antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a zinc finger protein which activates transcription of a scavenger receptor gene involved in the degradation of acetylated low density lipoprotein (Ac-LDL) (PMID: 11978792). This gene is located in a cluster of zinc finger genes on chromosome 19 at q13.4. A pseudogene of this gene is located on chromosome 15. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

55311

Swiss Prot

Q8N0Y2

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

EZF2; EZF-2; ZSCAN17; ZNF444

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using ZNF444 Rabbit pAb (A18470) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 90s.