# CD36/SR-B3 Rabbit mAb

Catalog No.: A19016 Recombinant 15 Publications



### **Basic Information**

**Observed MW** 80kDa

**Calculated MW** 53kDa

Category Primary antibody

Applications WB,ELISA

**Cross-Reactivity** Mouse, Rat

**CloneNo number** ARC0461

## Background

The protein encoded by this gene is the fourth major glycoprotein of the platelet surface and serves as a receptor for thrombospondin in platelets and various cell lines. Since thrombospondins are widely distributed proteins involved in a variety of adhesive processes, this protein may have important functions as a cell adhesion molecule. It binds to collagen, thrombospondin, anionic phospholipids and oxidized LDL. It directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes and it binds long chain fatty acids and may function in the transport and/or as a regulator of fatty acid transport. Mutations in this gene cause platelet glycoprotein deficiency. Multiple alternatively spliced transcript variants have been found for this gene.

### **Recommended Dilutions**

WB	1:1000 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
	requirements.

# Immunogen Information

Gene ID 948

**Swiss Prot** P16671

#### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

FAT; GP4; GP3B; GPIV; CHDS7; PASIV; SCARB3; BDPLT10; CD36/SR-B3

### Contact

6	400-999-6126
$\mathbf{X}$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn
€	www.abclonal.com.c

### **Product Information**


Source Rabbit

Isotype lgG

Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using CD36/SR-B3 Rabbit mAb (A19016) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1min.