

# NeuN Rabbit mAb

Catalog No.: A19086

Recombinant

21 Publications

## Basic Information

### Observed MW

46-55 kDa

### Calculated MW

34 kDa

### Category

Primary antibody

### Applications

WB,IF-F,IF-P,IHC-P,mIHC,ELISA

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

ARC0202

## Recommended Dilutions

<b>WB</b>	1:5000 - 1:20000
-----------	------------------

<b>IF-F</b>	1:200 - 1:800
-------------	---------------


<b>IF-P</b>	1:200 - 1:800
-------------	---------------

<b>IHC-P</b>	1:1000-1:4000
--------------	---------------

<b>mIHC</b>	1:1000-1:4000
-------------	---------------

<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
--------------	---

## Contact

	400-999-6126
--	--------------

	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
--	--

	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>
--	--

## Background

This gene encodes a member of the RNA-binding FOX protein family which is involved in the regulation of alternative splicing of pre-mRNA. The protein has an N-terminal proline-rich region, an RNA recognition motif (RRM) domain, and a C-terminal alanine-rich region. This gene produces the neuronal nuclei (NeuN) antigen that has been widely used as a marker for post-mitotic neurons. This gene has its highest expression in the central nervous system and plays a prominent role in neural tissue development and regulation of adult brain function. Mutations in this gene have been associated with numerous neurological disorders. Alternative splicing of this gene results in multiple transcript variants encoding distinct isoforms.

## Immunogen Information

### Gene ID

146713

### Swiss Prot

A6NFN3

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

FOX3; NEUN; FOX-3; HRNBP3; NeuN

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data




Western blot analysis of various lysates using NeuN Rabbit mAb (A19086) at 1:5000 dilution incubated overnight at 4°C.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Negative control (NC): C2C12, PC-12  
 Exposure time: 10s.




Confocal imaging of paraffin-embedded Human brain tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Mouse brain tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Rat brain tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



The multiplex IHC analysis on paraffin-embedded Mouse brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : NeuN Rabbit mAb (A19086, 1:2000) with TSA-TYR-520 (Green), GFAP Rabbit mAb (A19058, 1:500) with TSA-TYR-570 (Red), and TMEM119 Rabbit mAb (A27143, 1:600) with TSA-TYR-690 (Yellow). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M



The multiplex IHC analysis on paraffin-embedded Rat brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : NeuN Rabbit mAb (A19086, 1:2000) with TSA-TYR-520 (Green), GFAP Rabbit mAb (A19058, 1:500) with TSA-TYR-570 (Red), and TMEM119 Rabbit mAb (A27143, 1:600) with TSA-TYR-690 (Yellow). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M




Confocal imaging of frozen sections Mouse brain (Frozen section) tissue using NeuN Rabbit mAb (A19086, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

## Validation Data


---

citrate buffer at pH 6.0. The analysis was completed using a 20x objective lens.




Immunohistochemistry analysis of paraffin-embedded Human brain tissue using NeuN Rabbit mAb (A19086) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

citrate buffer at pH 6.0. The analysis was completed using a 20x objective lens.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using NeuN Rabbit mAb (A19086) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using NeuN Rabbit mAb (A19086) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.