

ICAM-1/CD54 Rabbit mAb

Catalog No.: A19300 **Recombinant** **5 Publications**

Basic Information

Observed MW

82-100kDa

Calculated MW

58kDa

Category

Primary antibody

Applications

WB,IHC-P,FC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC0261

Background

This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

FC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

3383

Swiss Prot

P05362

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

BB2; CD54; P3.58; MALA2; MyD10; ICAM-1/CD54

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of lysates from A549 cells using ICAM-1/CD54 Rabbit mAb (A19300) at 1:1000 dilution incubated overnight at 4°C. A549 cells were treated with TNF- α (20 ng/mL) at 37°C for 30 minutes.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 μ g per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 45s.




Flow cytometry: 1×10^6 293F cells (negative control, left) and Raji cells (right) were surface-stained with ICAM-1/CD54 Rabbit mAb (A19300, 10 μ g/mL, green line) or Rabbit IgG isotype control (AC042, 10 μ g/mL, blue line), followed by Alexa Fluor 647 conjugated goat anti-rabbit pAb (1:600 dilution) staining. Non-fluorescently stained cells were used as blank control (red line).




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human lung squamous carcinoma tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human lung tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Human pancreas tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using ICAM-1/CD54 Rabbit mAb (A19300) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.