

# SCAF8 Rabbit pAb

Catalog No.: A19467 **1 Publications**

## Basic Information

### Observed MW

170kDa

### Calculated MW

141kDa

### Category

Primary antibody

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

Enables RNA binding activity and RNA polymerase II C-terminal domain phosphoserine binding activity. Involved in negative regulation of termination of RNA polymerase II transcription, poly(A)-coupled and positive regulation of DNA-templated transcription, elongation. Located in nucleoplasm.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

22828

### Swiss Prot

Q9UPN6

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

RBM16; SCAF8

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG


### Purification

Affinity purification


### Storage

Store at -20°C

# Validation Data



Western blot analysis of various lysates using SCAF8 Rabbit pAb (A19467) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.



Western blot analysis of lysates from HeLa cells, using SCAF8 Rabbit pAb (A19467) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 180s.