[KD Validated] ERK1 Rabbit mAb

www.abclonal.com

ABclonal

Catalog No.: A19561 Recombinant 4 Publications

Basic Information

Observed MW

44kDa

Calculated MW

43kDa

Category

Primary antibody

Applications

WB,IP,IF/ICC,IF-P,IHC-P,ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC2591

Background

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described.

Recommended Dilutions

WB 1:1000 - 1:6000

0.5μg-4μg antibody for ΙP 200µg-400µg extracts of

whole cells

1:100 - 1:800 IF/ICC

IF-P 1:100 - 1:800

1:200 - 1:2000 **IHC-P**

Recommended starting **ELISA**

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 5595 P27361

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; K1

Product Information

Source Isotype **Purification** Rabbit IgG Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Contact

2	400-999-6126
\bowtie	cn.market@abclonal.com.cr
•	www.abclonal.com.cr



Immunoprecipitation analysis of 300 μg extracts of HeLa cells using 3 μg [KD Validated] ERK1 Rabbit mAb (A19561). Western blot was performed from the immunoprecipitate using ERK1 antibody (A19561) at a dilution of 1:1000.



Western blot analysis of lysates from wild type(WT) and ERK1 knockdown (KD) 293T cells, using [KD Validated] ERK1 Rabbit mAb (A19561) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of various lysates, using [KD Validated] ERK1 Rabbit mAb (A19561) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.




Confocal imaging of MCF7 cells using [KD Validated] ERK1 Rabbit mAb (A19561, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of paraffin-embedded 5xFAD mouse brain tissue using [KD Validated] ERK1 Rabbit mAb (A19561, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform microwave antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.



Immunohistochemistry analysis of paraffinembedded Human brain tissue using [KD Validated] ERK1 Rabbit mAb (A19561) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human breast tissue using [KD Validated] ERK1 Rabbit mAb (A19561) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human thyroid cancer tissue using [KD Validated] ERK1 Rabbit mAb (A19561) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.