

# STAT1 Rabbit mAb

Catalog No.: A19563   Recombinant   23 Publications

## Basic Information

### Observed MW

84 kDa/91 kDa

### Calculated MW

87 kDa

### Category

Primary antibody

### Applications

WB, IP, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Clone/No. number

ARC0042

## Background

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. The protein encoded by this gene can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. The protein plays an important role in immune responses to viral, fungal and mycobacterial pathogens. Mutations in this gene are associated with Immunodeficiency 31B, 31A, and 31C.

## Recommended Dilutions

|       |   |
|-------|---|
| WB    | 1:18000 - 1:60000   |
| IP    | 0.5 µg - 4 µg antibody for 200 µg-600 µg extracts of whole cells  |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |

## Immunogen Information

### Gene ID

6772

### Swiss Prot

P42224

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

CANDF7; IMD31A; IMD31B; IMD31C; ISGF-3; STAT91; STAT1

## Contact

|  |  |
|--|--|
|  | 400-999-6126   |
|  | <a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a> |
|  | <a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>             |

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data



Immunoprecipitation analysis of 600 µg extracts of HeLa cells using 3 µg STAT1 Rabbit mAb (A19563). Western blot was performed from the immunoprecipitate using STAT1 Rabbit mAb (A19563) at a dilution of 1:1000.



Western blot analysis of various lysates using STAT1 Rabbit mAb (A19563) at 1:20000 dilution incubated at room temperature for 1.5 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 20s.



Western blot analysis of various lysates using STAT1 Rabbit mAb (A19563) at 1:20000 dilution incubated at room temperature for 1.5 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 60s.