

[KO Validated] p53 Rabbit mAb

Catalog No.: A19585 **KO Validated** **Recombinant** **24 Publications**

Basic Information

Observed MW

53kDa

Calculated MW

44kDa

Category

Primary antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human

Clone/No. number

ARC56085

Background

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937277).

Recommended Dilutions

WB	1:10000 - 1:40000
IP	0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

7157

Swiss Prot

P04637

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

P53; BCC7; LFS1; BMFS5; TRP53; 53

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Product Information

Source	Isotype	Purification
Rabbit	IgG	Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Immunoprecipitation analysis of 300 μ g extracts from 293T cells using 3 μ g [KO Validated] p53 Rabbit mAb (A19585). Western blot was performed from the immunoprecipitate using [KO Validated] p53 Rabbit mAb (A19585) at a dilution of 1:1000.



Western blot analysis of lysates from wild type (WT) and p53 knockout (KO) 293T cells using [KO Validated] p53 Rabbit mAb (A19585) at 1:10000 dilution incubated at room temperature for 1.5 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.