# [KO Validated] TIMM17A Rabbit pAb

www.abclonal.com

**ABclonal** 

Catalog No.: A19917 KO Validated

### **Basic Information**

### **Observed MW**

16kDa

### **Calculated MW**

18kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IF/ICC

### **Cross-Reactivity**

Human, Mouse

## **Background**

Predicted to contribute to protein transmembrane transporter activity. Predicted to be involved in protein import into mitochondrial matrix. Located in mitochondrial inner membrane and nucleoplasm. Is integral component of mitochondrial inner membrane. Part of TIM23 mitochondrial import inner membrane translocase complex.

## **Recommended Dilutions**

1:500 - 1:2000 **WB** 

1:50 - 1:200 IF/ICC

## **Immunogen Information**

**Gene ID Swiss Prot** 10440 Q99595

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-171 of human TIMM17A (NP\_006326.1).

## **Synonyms**

TIM17; TIM17A; 7A

### **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn

### **Product Information**


**Purification Source** Isotype Rabbit IgG Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from wild type (WT) and TIMM17A knockout (KO) 293T cells, using [KO Validated] TIMM17A Rabbit pAb (A19917) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.