

Somatostatin (SST) Rabbit mAb

Catalog No.: A20617 **Recombinant** **1 Publications**

Basic Information

Observed MW

Refer to figures

Calculated MW

13kDa

Category

Primary antibody

Applications

ELISA, IHC-P, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC50670

Background

The hormone somatostatin has active 14 aa and 28 aa forms that are produced by alternate cleavage of the single preproprotein encoded by this gene. Somatostatin is expressed throughout the body and inhibits the release of numerous secondary hormones by binding to high-affinity G-protein-coupled somatostatin receptors. This hormone is an important regulator of the endocrine system through its interactions with pituitary growth hormone, thyroid stimulating hormone, and most hormones of the gastrointestinal tract. Somatostatin also affects rates of neurotransmission in the central nervous system and proliferation of both normal and tumorigenic cells.

Recommended Dilutions

IHC-P 1:100 - 1:500

IF/ICC 1:50-1:200

Immunogen Information

Gene ID

6750

Swiss Prot

P61278

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 23-116 of human Somatostatin (SST) (NP_001039.1).

Synonyms

SMST; SST1; Somatostatin (SST)

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.



Immunohistochemistry analysis of Somatostatin (SST) in paraffin-embedded human pancreas tissue using Somatostatin (SST) Rabbit mAb (A20617) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Somatostatin (SST) in paraffin-embedded mouse pancreas tissue using Somatostatin (SST) Rabbit mAb (A20617) at a dilution of 1:500 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Human pancreas tissue using Somatostatin (SST) Rabbit mAb (A20617, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.