# **CPT1A Rabbit mAb**

Catalog No.: A20746 Recombinant 16 Publications



# **Basic Information**

#### **Observed MW**

88kDa

#### **Calculated MW**

88kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human

#### CloneNo number

ARC51171

# **Background**

The mitochondrial oxidation of long-chain fatty acids is initiated by the sequential action of carnitine palmitoyltransferase I (which is located in the outer membrane and is detergent-labile) and carnitine palmitoyltransferase II (which is located in the inner membrane and is detergent-stable), together with a carnitine-acylcarnitine translocase. CPT I is the key enzyme in the carnitine-dependent transport across the mitochondrial inner membrane and its deficiency results in a decreased rate of fatty acid beta-oxidation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

WB 1:1000 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**Swiss Prot
1374
P50416

# Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

CPT1; CPT1-L; L-CPT1; CPT1A

### **Contact**

<b>a</b>	400-999-6126
$\sim$	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Western blot analysis of lysates from MCF7 cells, using CPT1A Rabbit mAb (A20746) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.