

# E-Cadherin Rabbit mAb

Catalog No.: A20798 **Recombinant** **107 Publications**

## Basic Information

### Observed MW

125 kDa

### Calculated MW

97 kDa

### Category

Primary antibody

### Applications

WB,IF/ICC,IF-F,IF-P,IHC-P,ELISA

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

ARC51012

## Background

This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function of this gene is thought to contribute to cancer progression by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. This gene is present in a gene cluster with other members of the cadherin family on chromosome 16.

## Recommended Dilutions

**WB** 1:1000 - 1:2000

**IF/ICC** 1:100 - 1:1000

**IF-F** 1:200 - 1:600

**IF-P** 1:100 - 1:1000

**IHC-P** 1:400 - 1:2000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

999

### Swiss Prot

P12830

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

UVO; CDHE; ECAD; LCAM; Arc-1; BCDS1; CD324; E-Cadherin

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Contact

☎ | 400-999-6126


✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)


## Validation Data



Western blot analysis of various lysates, using E-Cadherin Rabbit mAb (A20798) at 1:1000 dilution incubated overnight at 4°C.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Negative control (NC): HeLa  
Exposure time: 30s.




Western blot analysis of lysates from Rat kidney using E-Cadherin Rabbit mAb (A20798) at 1:1000 dilution incubated overnight at 4°C.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 180s.




Western blot analysis of lysates from Mouse lung using E-Cadherin Rabbit mAb (A20798) at 1:2000 dilution incubated overnight at 4°C.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 45s.


## Validation Data




Confocal imaging of paraffin-embedded Human prostate cancer tissue using E-Cadherin Rabbit mAb (A20798, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Mouse pancreas tissue using E-Cadherin Rabbit mAb (A20798, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of MCF7 cells using E-Cadherin Rabbit mAb (A20798, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with  $\alpha$ -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of paraffin-embedded Human colon tissue using E-Cadherin Rabbit mAb (A20798, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen




Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen



Immunohistochemistry analysis of paraffin-embedded Human pancreas tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen


## Validation Data

retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human placenta tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.

retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.


retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse intestin tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using E-Cadherin Rabbit mAb (A20798) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.



Confocal imaging of frozen sections of Mouse colon tissue using E-Cadherin Rabbit mAb (A20798, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Rat colon tissue using E-Cadherin Rabbit mAb (A20798, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500)

## Validation Data

---

(Red). DAPI was used for nuclear staining  
(Blue). Microwave antigen retrieval  
performed with 0.01M Citrate Buffer (pH 6.0)  
prior to IF staining. Objective: 40x.