# **GNA15 Rabbit pAb**

Catalog No.: A2080



## **Basic Information**

#### **Observed MW**

43kDa

### **Calculated MW**

44kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Mouse, Rat

## **Background**

Enables G protein-coupled receptor binding activity. Involved in positive regulation of cytosolic calcium ion concentration involved in phospholipase C-activating G protein-coupled signaling pathway. Predicted to be located in plasma membrane. Predicted to be part of heterotrimeric G-protein complex.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting

concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**2769

Swiss Prot
P30679

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

HG1L; GNA16; GNA15

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{\triangle}$	ī	www.ahclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of various lysates using GNA15 Rabbit pAb (A2080) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.