# IRE1 Rabbit mAb

Catalog No.: A21021 Recombinant 9 Publications



# **Basic Information**

### **Observed MW**

130kDa

### **Calculated MW**

110kDa

## Category

Primary antibody

### **Applications**

WB,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC57190

## **Background**

This gene encodes the transmembrane protein kinase inositol-requiring enzyme 1. The encoded protein contains two functional catalytic domains, a serine/threonine-protein kinase domain and an endoribonuclease domain. This protein functions as a sensor of unfolded proteins in the endoplasmic reticulum (ER) and triggers an intracellular signaling pathway termed the unfolded protein response (UPR). The UPR is an ER stress response that is conserved from yeast to mammals and activates genes involved in degrading misfolded proteins, regulating protein synthesis and activating molecular chaperones. This protein specifically mediates the splicing and activation of the stress response transcription factor X-box binding protein 1.

# **Recommended Dilutions**

**WB** 1:2000 - 1:6500

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID**2081

Swiss Prot
075460

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

IRE1; IRE1P; IRE1a; hIRE1p

## **Contact**

| <b>a</b>  |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| •         | ī | www.abclonal.com.cn       |

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of various lysates using IRE1 Rabbit mAb (A21021) at 1:6000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.